

# COMMISSIONED REVIEW ARTICLE

## Exercise acts as a drug; the pharmacological benefits of exercise

J Vina, F Sanchis-Gomar, V Martinez-Bello and MC Gomez-Cabrera

*Department of Physiology, University of Valencia, Fundacion Investigacion Hospital Clinico  
Universitario/INCLIVA, Valencia, Spain*

### Correspondence

Professor Jose Viña, Department  
of Physiology, Faculty of  
Medicine, University of Valencia,  
Av. Blasco Ibañez, 15, Valencia,  
46010 Spain. E-mail:  
jose.vina@uv.es

### Keywords

health; dosing of exercise;  
contraindications of exercise;  
sport; training

### Received

25 January 2012

### Revised

11 March 2012

### Accepted

16 March 2012

The beneficial effects of regular exercise for the promotion of health and cure of diseases have been clearly shown. In this review, we would like to postulate the idea that exercise can be considered as a drug. Exercise causes a myriad of beneficial effects for health, including the promotion of health and lifespan, and these are reviewed in the first section of this paper. Then we deal with the dosing of exercise. As with many drugs, dosing is extremely important to get the beneficial effects of exercise. To this end, the organism adapts to exercise. We review the molecular signalling pathways involved in these adaptations because understanding them is of great importance to be able to prescribe exercise in an appropriate manner. Special attention must be paid to the psychological effects of exercise. These are so powerful that we would like to propose that exercise may be considered as a psychoactive drug. In moderate doses, it causes very pronounced relaxing effects on the majority of the population, but some persons may even become addicted to exercise. Finally, there may be some contraindications to exercise that arise when people are severely ill, and these are described in the final section of the review. Our general conclusion is that exercise is so effective that it should be considered as a drug, but that more attention should be paid to the dosing and to individual variations between patients.

### Abbreviations

ACSM, American College of Sports Medicine; ACTH, adrenocorticotrophic hormone; AICAR, 5-aminoimidazole-4-carboxamide-1- $\beta$ -D-ribofuranoside; AMPK, AMP-activated protein kinase; BDNF, brain-derived neurotrophic factor; CHD, coronary heart disease; GDNF, glial cell-derived neurotrophic factor; HIF-1, hypoxia-inducible factor 1; HSF, heat shock factor; HSP, heat shock protein; IGF, insulin growth factor; MET, metabolic equivalent (estimated oxygen cost of 3.5 mL·min<sup>-1</sup>·kg<sup>-1</sup>); MnSOD, mitochondrial superoxide dismutase; PGC-1 $\alpha$ , PPAR- $\gamma$  coactivator-1 $\alpha$ ; ROS, reactive oxygen species; SCD, sudden cardiac death; VO<sub>2</sub>, oxygen consumption

### Exercise, movement and health: definitions

Health promotion is the science and art of helping people change their lifestyle to move towards a state of optimal health (O'Donnell, 1986). The World Health Organization defines health as 'Physical, mental, and social well-being, not merely the absence of disease and infirmity'. Physical fitness is defined as the physiologic state of well-being that allows one to meet the demands of daily living (health-related physical fitness) or that provides the basis for sport performance (performance-related physical fitness), or both. Although we are aware that there is a clear difference between

the terms physical activity ('any bodily movement') and exercise ('a subset of physical activity that is characterized by a planned and purposeful training') (Caspersen *et al.*, 1985), in this review, we are going to use these two concepts as synonymous because some of the studies to which we will refer use the terms interchangeably.

### Historical background

The hypothesis that physical activity promotes health and longevity is not new. As far back as 2500 BC, in ancient China, records of organized exercise for health promotion

1 [?] have been found (Lyons and Rjja, 1978; Lee and Skerrett,  
2 2001). In Greco-Roman times, 2500 years ago, Hippocrates  
3 (460–370 BC) and later Galen (AD 129–210) recognized the  
4 need to promote and prescribe exercise for health-related  
5 benefits and the need to provide general medical care for the  
6 athletic individual (Speed and Jaques, 2010). In this regard,  
7 the philosopher Plato (427–347 BC) said: ‘Lack of activity  
8 destroys the good condition of every human being while  
9 movement and methodical physical exercise saves and pre-  
10 serves it’ (Fox and Haskell, 1968).

11 Simple comparisons of men in different occupations pro-  
12 vided the first empirical evidence that physical activity was  
13 associated with health. The first studies demonstrating a sig-  
14 nificant inverse relationship between physical activity and  
15 coronary heart disease (CHD) were those conducted by  
16 Morris *et al.* (1953b) in London early in the 1950s. These  
17 authors found that London bus conductors had only 73% the  
18 frequency of CHD that was found in the less active bus  
19 drivers. Their later comparison of London postmen and less  
20 active postal clerks produced much the same findings (Morris  
21 *et al.*, 1953a). These seminal studies were followed by those of  
22 Paffenbarger and collaborators in the 1970s, assessing the  
23 increase in the relative risk of death from any cause and from  
24 specific diseases associated with physical inactivity (Paffen-  
25 barger and Hale, 1975; Paffenbarger *et al.*, 1978).

## 27 Exercise is beneficial for your health

28 Exercise is one of the most frequently prescribed therapies  
29 both in health and disease. There is irrefutable evidence  
30 showing the beneficial effects of exercise both to prevent  
31 and to treat several diseases. Researchers have shown that  
32 both men and women who report increased levels of phys-  
33 ical activity and fitness have reductions in relative risk of  
34 death (by about 20%–35%) (Blair *et al.*, 1989; Macera *et al.*,  
35 2003).

36 Recent research suggests that modest increments in  
37 energy expenditure due to physical activity (~1000 kcal per  
38 week) or an increase in physical fitness of 1 MET (metabolic  
39 equivalent) is associated with lowering mortality by about  
40 20% (Myers *et al.*, 2004). Physically inactive middle-aged  
41 women (engaging in less than 1 h of exercise per week) expe-  
42 rience a 52% increase in all-cause mortality, a doubling of  
43 cardiovascular-related mortality, and a 29% increase in  
44 cancer-related mortality when compared with physically  
45 active ones (Hu *et al.*, 2004). Thus, there is clear evidence that  
46 regular physical activity produces significant health effects  
47 and reduces the risk of premature death from any cause and  
48 from cardiovascular disease in particular amongst asymptom-  
49 atic men and women.

50 The benefits of physical activity are evident, not only in  
51 healthy persons but also in patients. Observational and ran-  
52 domized trials have shown that regular physical activity con-  
53 tributes to the treatment of several chronic diseases  
54 (Bouchard *et al.*, 1994; Warburton *et al.*, 2006a). There is evi-  
55 dence for prescribing exercise in the primary and secondary  
56 prevention of pulmonary and cardiovascular diseases (CHD,  
57 chronic obstructive pulmonary disease, hypertension, inter-  
58 mittent claudication); metabolic disorders (type 2 diabetes,  
59 dyslipaemia, obesity, insulin resistance); muscle, bone and

60 joint diseases (rheumatoid arthritis, fibromyalgia, chronic  
61 fatigue syndrome, osteoporosis); cancer; and depression (Ped-  
62 ersen and Saltin, 2006; Warburton *et al.*, 2006a). Even if exer-  
63 cise is an effective therapeutic agent for all of these diseases,  
64 as with any other medicine, the dosage (volume and intensity  
65 of the exercise), frequency of administration (sessions per  
66 week), type (aerobic vs. resistance exercise), systemic and  
67 psychoactive effects and contraindications and side effects of  
68 the exercise must be taken into account to achieve the best  
69 clinical outcome. For instance, both resistance and aerobic  
70 training have been shown to be of benefit for the control of  
71 diabetes; however, resistance training may have greater ben-  
72 efits for glycaemic control than aerobic training (Dunstan  
73 *et al.*, 2005).  
74

## 75 The dosage of exercise

76 Dosage is important in clinical medicine and all marketed  
77 drugs require data on their efficacy and safety (Lee, 2007). It  
78 is known that there is a minimum amount of physical activ-  
79 ity for health benefits. These benefits increase with increasing  
80 the amount of exercise, but beyond a certain level, adverse  
81 effects outweigh benefits (Lee, 2007). Unlike chemical drugs,  
82 however, the minimum dose, dose response and maximum  
83 safe dose of physical activity are not well understood (Lee,  
84 2007). There is a continuous debate on how much, what type,  
85 how often, what intensity and how lengthy physical activity  
86 should be. This is important for issuing public health recom-  
87 mendations (Blair *et al.*, 2004). Summarizing available infor-  
88 mation across studies is difficult because investigators have  
89 measured exercise intensity in different ways and classified  
90 physical activity according to different dose schemes that are  
91 often difficult to compare (Lee, 2007). Over the years, various  
92 expert groups, based on the best evidence available, have  
93 postulated different physical activity recommendations and  
94 guidelines (see Table 1).  
95

96 Intensity levels of physical activity can be expressed rela-  
97 tive to oxygen consumption ( $VO_2$ ) or to heart rate (Warburton  
98 *et al.*, 2006b). Moderate-intensity activities are those in  
99 which heart rate and breathing are raised; but, still, it is  
100 possible to speak comfortably. This occurs around 4–6 METs  
101 and brisk walking at 3.0 mph (80.4 m·min<sup>-1</sup>) is one such  
102 activity. Vigorous-intensity activities are that in which heart  
103 rate is higher, breathing is heavier and conversation is harder  
104 (about 6–8 METs) (Warburton *et al.*, 2006b), for instance  
105 jogging. It has been shown that exercising at even 50% of the  
106 recommended levels (72 min of moderate exercise a week)  
107 appears sufficient to provide some improvement in fitness.  
108 However, at this low exercise dosage, cardiovascular risk  
109 factors (blood pressure, lipid profile and weight) do not  
110 improve (Church *et al.*, 2007). In fact, for many individuals,  
111 up to 60 min of daily physical activity are more appropriate if  
112 weight control is the primary goal (Lee, 2007). Thus, dose-  
113 response relations between physical activity and different  
114 health outcomes are different. The evaluation of the  
115 minimum amount of physical activity (lower dose) necessary  
116 to achieve its beneficial effects has been the object of intense  
117 research. Wen *et al.* (2011) have recently found that 15 min a  
118 day or 90 min a week of moderate-intensity exercise is of  
119

**Table 1**

Historical evolution in physical activity recommendations and guidelines

	Physical activity recommendations	Intensity	Minutes	Frequency	Reference
1970s–1980s	Vigorous exercise (e.g. running)		20 min·day <sup>-1</sup>	3 times·week <sup>-1</sup>	(1978)
1990s	Moderate exercise (e.g. brisk walking)		30 min·day <sup>-1</sup>	Most days of the week	(Pate <i>et al.</i> , 1995; 1996)
2000s	Moderate exercise		60 min·day <sup>-1</sup>	3 times·week <sup>-1</sup>	(Lee, 2007)
2010 (healthy adults ages 18–45)	Moderate exercise		30 min·day <sup>-1</sup> (150 min week <sup>-1</sup> )	Most days of the week (5 days·week <sup>-1</sup> )	(O'Donovan <i>et al.</i> , 2010)
	Vigorous exercise		75 min·week <sup>-1</sup>		(O'Donovan <i>et al.</i> , 2010)

benefit in terms of life expectancy, even for subjects with cardiovascular risks.


### Exhaustive exercise and longevity

Although the health benefits of leisure-time physical activity are well documented, the association between vigorous exercise training and mortality or longevity of elite athletes is not fully understood (Teramoto and Bungum, 2010). For centuries, the general belief has been that exhaustive, competitive exercise is harmful and decreases life expectancy (Ruiz *et al.*, 2010). For instance, Moorstein (1968) stated that all members of the 1948 Harvard rowing crew had died early from cardiac diseases. In contrast, it has been shown that participation in endurance competitive sports increases life expectancy. It was found that the life expectancy of oarsmen was higher than that of their non-athletic controls (Hartley and Llewellyn, 1939; Prout, 1972). Karvonen and co-workers found that Finnish champion skiers (born between 1845 and 1910) lived 2.8–4.3 years longer than general male population in Finland. In contrast with most studies of that time, Polednak (1972) reported evidence against the beneficial effects of strenuous exercise. He found differences in longevity and cardiovascular mortality related to the extent of participation in college athletics. Moreover, in a recent animal study, it has been found that long-term vigorous endurance exercise training may in some cases promote adverse cardiac remodelling and produce a substrate for cardiac arrhythmias (Benito *et al.*, 2011). The incidence of sudden cardiac death (SCD) amongst young athletes (estimated to be 1–3 per 100 000 person-years) is higher than in non-athletes and may possibly still be underestimated (Drezner, 2008). However, it has been shown that the most common cause of SCD in young athletes is underlying inherited cardiac disease, such as cardiomyopathies, congenital coronary anomalies and ion channelopathies (Maron *et al.*, 2009). To clarify this apparent contradiction, we determined the longevity of the participants of the Tour de France and compared it with that of the general population born between 1892 and 1942. The Tour de France is amongst the most gruelling sport events in the world. We found an 11% increase in average longevity in Tour de France participants when compared with the general

population (Sanchis-Gomar *et al.*, 2011b) (see Figure 1). Thus, the majority of data in human studies support the notion that prescription of regular, vigorous aerobic exercise might be a useful tool, with a dose–effect response to improve the overall health status and longevity of the general population (Ruiz *et al.*, 2010; Teramoto and Bungum, 2010). In our opinion, physicians, health professionals and general population should not be under the impression that strenuous exercise and/or high-level aerobic competitive sports are bad for one's health and shorten one's life. Thus, a dose–response relation appears to exist, such that people who have the highest levels of physical activity and fitness are at lowest risk of premature death (Warburton *et al.*, 2006a).

Training status is a very relevant factor in the prescription of the exercise 'dose'. Increasing the doses of exercise has positive consequences for health in trained individuals (Ruiz *et al.*, 2010; Sanchis-Gomar *et al.*, 2011b), whereas heavy physical exertion can trigger the onset of acute myocardial infarction, particularly in people who are habitually sedentary (Mittleman *et al.* 1993). Results from the same group showed that less active men participating in vigorous activity were more likely to have a myocardial infarction during exercise than the most active men (Thompson *et al.*, 2007).

In the pharmacological treatment of many conditions, physicians typically start with a dose of a drug believed to be the minimum effective dose. If the patient does not respond, this initial dose may then be titrated upwards to a maximum dose, beyond which the adverse effects of the drug are unacceptable for treatment (Lee, 2007). Thus, the intensity of aerobic training may be also titrated in healthy people (Warburton *et al.*, 2006b). Unfit people can get significant improvements in physical fitness with a low training intensity, while those with a higher fitness level need a greater level of exercise intensity to achieve further improvements in fitness (Shephard, 2001). Thus, these fit individuals who have met the physical activity levels recommended for all healthy adults for at least 6 months may obtain additional health benefits by engaging in 300 min or more of moderate-intensity aerobic activity per week, or 150 min or more of vigorous-intensity aerobic activity each week, or equivalent combinations of moderate- and vigorous-intensity aerobic activities (Lee and Skerrett, 2001; O'Donovan *et al.*, 2010). These relatively low doses are, obviously, not applicable to



**Figure 1**

Percentage of survival related to age in Tour de France (Tdf) participants and in the general population. Persons born between 1892 and 1942 have been studied. Average life span of Tour de France participants is higher ( $P = 0.004$ ; 17.5%) than the general population of the same country in which the cyclists were born. The age at which 50% of the general population died was 73.5 years, compared with 81.5 years in Tour de France participants (i.e. 11% increase).

high level professional athletes who perform exercise at much higher doses.

The guidelines discussed above are generally appropriate for young to middle-aged adults. But, as with medicines, special considerations should be taken when prescribing exercise for people with special needs such as elderly, children, pregnant women, overweight or obese patients and patients with chronic diseases (Warburton *et al.*, 2006b). For instance, it has been shown that vigorous activities are not essential for the reduction of cardiovascular risk in men over 60. Regular physical activity is enough to achieve a significant decrease in mortality in this population. Thus, the greatest benefit to health is gained from sustained moderate exercise, above which there appears to be no further benefit to health in older men (Hakim *et al.*, 1998; Wannamethee *et al.*, 1998).


Regarding the 'dosage' of exercise, whether it should be performed in either one continuous or two or more accumulated bouts, the available evidence suggest that at least for fitness, accumulated and continuous patterns of exercise training of the same total duration confer similar benefits (Murphy *et al.*, 2009). For instance, it has been shown that five to eight 2 min bouts of stair climbing accumulated over the course of a day confer health benefits, including increases in cardiovascular fitness, compared with non-exercising controls (Boreham *et al.*, 2005).

Although physical activity is beneficial to health with or without weight loss, adults who find it difficult to maintain a normal weight and adults with increased risk of cardiovascular disease or type 2 diabetes, in particular, may benefit from going beyond the levels of activity recommended for all healthy adults and gradually progressing towards meeting the recommendations for conditioned individuals (O'Donovan *et al.*, 2010).

## Systemic adaptations to exercise

The exercise-induced adaptations are especially evident in the cardiorespiratory and musculoskeletal systems, and body composition and metabolism (Warburton *et al.*, 2006a; Lee *et al.*, 2010). But the documented health benefits of exercise also include diminished symptoms of depression and anxiety (Kujala, 2011).

Skeletal muscle is the main target of exercise training. Modifications in skeletal muscle are crucial for enhancing endurance and metabolic efficiency (Matsakas and Narkar, 2010). Muscle fibres are commonly classified as type I slow-twitch or oxidative fibres, with a high mitochondrial content, and type II fast-twitch or glycolytic fibres, which have fewer mitochondria. Endurance exercise induces an increase in



**Figure 2**

Health benefits of exercise in tissues and organs.

mitochondriogenesis, a shift in fibre distribution from glycolytic to oxidative and an increase in fatty acid oxidation that ultimately leads to an increase in aerobic capacity and retards diseases such as obesity, type 2 diabetes and cardiovascular diseases (Holloszy and Coyle, 1984; Mootha *et al.*, 2003).

It has been shown that regular exercise can reduce abdominal adiposity and improve weight control (Warburton *et al.*, 2006a), enhance lipoprotein profiles (e.g. reduce triglyceride levels, increase high density lipoprotein and decrease low-density lipoprotein levels), improve glucose homeostasis and insulin sensitivity, reduce blood pressure, improve autonomic tone, reduce systemic inflammation; decrease blood coagulation, improve coronary blood flow, augment cardiac function and enhance endothelial function (Warburton *et al.*, 2006a).

Regular physical activity is also associated with improved psychological well-being (e.g. through reduced stress, anxiety and depression) (Dunn *et al.*, 2001). The beneficial effects of exercise on cognitive function are well known (Neeper *et al.*, 1995). The mechanism behind this is not fully understood, but it seems to be associated with an increased expression of neurotrophic factors in some brain areas. Increased expression of these factors is related to better memory and improved cognitive function. Brain-derived neurotrophic factor (BDNF) can enhance the survival and differentiation of neurons, and voluntary exercise has been shown to increase it (Neeper *et al.*, 1996). Psychological well-being is particularly important for the prevention and management of cardiovascular disease, but it also has important implications for the prevention and management of other chronic diseases such as diabetes, osteoporosis, hypertension, obesity, cancer and depression (Warburton *et al.*, 2006a). It has been shown that physical activity results in specific adaptations that affect individual states in all of these diseases. For instance, adaptations that affect glucose homeostasis, in type 2 diabetes, are of great importance. Several changes occur as a result of


regular physical activity, including increased glycogen synthase and hexokinase activities, increased mRNA and protein expression of the glucose transporter GLUT-4 and improved muscle capillary density (resulting improved glucose delivery to the muscle) (Mandroukas *et al.*, 1984).

Exercise causes a significant reduction in cancer rates (specifically colon and breast cancer) (Shephard and Futcher, 1997; Pedersen and Saltin, 2006). Possible explanations include reductions in fat stores, increased energy expenditure offsetting a high fat diet, activity-related changes in sex hormone levels, immune function, insulin and insulin-like growth factors, free radical generation and direct effects on the tumour cell biology (Westerlind, 2003).

The majority of the proposed mechanisms have been discussed in the context of chronic adaptations by regular physical activity. However, it has been shown that isolated exercise sessions (separate doses of exercise) also elicit transient, but still beneficial, changes in risk factors for chronic diseases (Thompson *et al.*, 2001). Many of the training adaptations derive from a single exercise bout that elicits cellular changes at the gene level leading to cumulative effects of training. The acute effect of exercise results in transient reductions in triglyceride levels, increases in HDL cholesterol level, decreases in blood pressure, reductions in insulin resistance and improvements in glucose control (Thompson *et al.*, 2001). These acute changes underpin the important role that individual exercise sessions have on health status. Thus, single doses of exercise have also a relevant impact on health. Figure 2 summarizes the favourable effects of exercise.

### Signalling pathways regulated by exercise in skeletal muscle

The regulation of the cellular functions with exercise are dependent on many stimuli: alterations in metabolite con-



**Figure 3**

A summary of the signalling pathways regulated by exercise in skeletal muscle.

centrations, a shift in the ATP : ADP ratio, changes in the intracellular concentration of  $Ca^{2+}$ , in the intracellular pH, and activations of the oxidative stress-sensitive signalling pathways (Sakamoto and Goodyear, 2002; Ji *et al.*, 2004). To elucidate the molecular signalling mechanisms that enable skeletal muscle to respond to the contractile stimulus and that mediate the adaptations to exercise is of great importance (Sakamoto and Goodyear, 2002). It has been clearly established that physical exercise can activate MAPK signalling, including the ERK1/2 (Goodyear *et al.*, 1996), p38 (Gomez-Cabrera *et al.*, 2005) and JNK pathways (Goodyear *et al.*, 1996). It can also increase the activity of the AMP-activated protein kinase (AMPK), Akt and the p70 S6 kinase (Sakamoto and Goodyear, 2002). In skeletal muscle  $Ca^{2+}$  signalling is extensive. In addition to triggering muscle contraction through the troponin system,  $Ca^{2+}$  is also involved in the regulation of relevant intracellular proteins such as PKC, calcineurin and calmodulin kinase that mediate cellular signal transduction (Berchtold *et al.*, 2000).

More recently, it has been shown, that low-to-moderate levels of reactive oxygen species (ROS) play multiple regulatory roles in cells such as the control of gene expression, regulation of cell signalling pathways and modulation of skeletal muscle force production (Reid, 2001) (see Figure 3).

### ROS and exercise: training as an antioxidant intervention

The role of ROS in the exercise-induced adaptations in skeletal muscle has been extensively studied (Salminen and Vihko, 1983; Gomez-Cabrera *et al.*, 2008b). The idea of the deleterious effects of ROS has been firmly entrenched in the minds of scientists during the last 30 years. However, there is growing evidence that the continued presence of low con-

centrations of free radicals is, in fact, able to induce the expression of antioxidant enzymes and other defence mechanisms. In this scenario, radicals may be seen as beneficial as they act as signals to enhance defences rather than deleterious as they are when cells are exposed to high levels of these radicals. Animals frequently exposed to chronic exercise have shown less oxidative damage after exhaustive exercise than untrained ones (Salminen and Vihko, 1983). This is largely due to the up-regulation of endogenous antioxidant enzymes such as glutathione peroxidase, mitochondrial superoxide dismutase (MnSOD) and  $\gamma$ -glutamylcysteine synthetase (Salminen and Vihko, 1983). A major conclusion that can be drawn from these results is that exercise itself acts as an antioxidant, because training increases the expression of antioxidant enzymes (Gomez-Cabrera *et al.*, 2008b). Subsequently, we and others have shown that antioxidant supplements prevent the induction of mitochondrial biogenesis, molecular regulators of insulin sensitivity and endogenous antioxidant defence by physical exercise (Gomez-Cabrera *et al.*, 2008a; Ristow *et al.*, 2009). Thus, ROS act as signals in exercise because decreasing their formation prevents activation of important signalling pathways, which cause useful adaptations in cells.

Because of the widespread implications of ROS in almost all important biological functions, it is difficult to define all the pathways and gene targets that are affected by redox signalling during exercise. The following are some of the most relevant signalling pathways modulated by exercise: the PPAR- $\gamma$  coactivator-1 $\alpha$  and  $\beta$  (PGC-1 $\alpha$  and PGC-1 $\beta$ ) (Gomez-Cabrera *et al.*, 2008a; Ristow *et al.*, 2009), p53 (Borras *et al.*, 2011), hypoxia-inducible factor 1 (HIF-1) (Huang *et al.*, 1996), heat shock factor (HSF) (Palomero *et al.*, 2008), NF- $\kappa$ B and MAPK signalling pathways (Ji *et al.*, 2004; Gomez-Cabrera *et al.*, 2005). Several important adaptations in skeletal muscle such as mitochondrial biogenesis, antioxidant defence, hypertrophy, cytoprotection and fibre transforma-

tion are regulated primarily by these pathways. Thus, its regulation should be tightly controlled (Gomez-Cabrera *et al.*, 2009) (see Figure 3).

## Exercise, a psychoactive drug

The effects of exercise training on brain function have received much attention. In the early '80s, exercise was shown to increase  $\beta$ -endorphin in peripheral blood in humans (Bortz *et al.*, 1981; Carr *et al.*, 1981). Elevated serum  $\beta$ -endorphin concentrations induced by exercise have since been linked to a variety of psychological and physiological changes, including mood state changes and 'exercise-induced euphoria', altered pain perception and responses to numerous stress hormones (growth hormone, ACTH, prolactin, catecholamines and cortisol) (Harber and Sutton, 1984).

Exercise training can favourably influence cognitive function (Dishman *et al.*, 2006; Vaynman and Gomez-Pinilla, 2006). Exercise improves learning and memory (van Praag *et al.*, 1999), improves the quality of sleep, counteracts the mental decline that comes with age (Laurin *et al.*, 2001) and facilitates functional recovery from brain injury (Grealy *et al.*, 1999) and depression (Siuciak *et al.*, 1996; Shirayama *et al.*, 2002). Exercise is a very powerful stimulus to the induction of neurogenesis in the adult dentate gyrus (van Praag *et al.*, 1999) that can contribute to remodelling hippocampal synaptic circuits and to enhance cognitive function.

Exercise training can also mitigate the consequences of acute exposure to different types of psychological stress (Dishman *et al.*, 2006) and exercise-induced alterations in the 5-hydroxytryptaminergic and the noradrenergic systems can explain these responses (Dishman *et al.*, 2006). Most of the positive effects of exercise, as mentioned previously, have been related to the induction, in different brain areas, of neurotrophic proteins, including BDNF, glial cell-derived neurotrophic factor (GDNF) and insulin growth factor (IGF). Whether brain metabolic responses to acute physical activity extend beyond regions specifically involved with motor, sensory or cardiovascular autonomic control is not as yet clear (Dishman *et al.*, 2006). Transient increases in local cerebral glucose use and in cerebral blood flow have been reported in the different brain areas in response to acute strenuous treadmill running in rats and in humans (Vissing *et al.*, 1996). Also, the discharge rate of a select pool of hippocampal pyramidal cells increased as running velocity increased (Czurko *et al.*, 1999). Moreover, exercise increased metabolic capacity in the motor cortex and striatum (McCloskey *et al.*, 2001).

The psychoactive effects of exercise that we have just mentioned are not free from risks. Pathological patterns of behaviour in gym clients have been reported (Lejoyeux *et al.*, 2008). As observed in patients with eating disorders, active individuals usually worry about their body shape, put special attention on their eating patterns, show exercise addiction and have a perfectionism personality trait (Freimuth *et al.*, 2011). This body image disorder has been addressed as reverse anorexia, vigorexia or muscle dysmorphia (Lejoyeux *et al.*, 2008). Based on a review of a wide range of studies on exercise addiction, it has been estimated that its prevalence in the general population is close to 3%. Amongst certain groups

such as ultra-marathon runners, body builders and sport science students, the percentage is even higher (Freimuth *et al.*, 2011; Sussman and Sussman, 2011).

## Contraindications for exercise

The purpose of this section is to discuss why under some circumstances physical exercise does not increase the quality of life.

Although both the heart and the lung benefit significantly from physical activity, there are some contraindications when exercise is performed by patients suffering from heart and pulmonary diseases. Pedersen and Saltin (2006) reviewed the possible contraindications of exercise in most of the diseases in which exercise have shown beneficial effects. For instance in patients with CHD, exercise is contraindicated until the condition has been stable for at least 5 days; dyspnea at rest, aortic stenosis, pericarditis, myocarditis, endocarditis, fever and severe hypertension all are contraindications to exercise (Pedersen and Saltin, 2006). Black *et al.* (1975) were amongst the first to find that strenuous exercise can cause acute injury to coronary plaques, leading to occlusion of coronary arteries. However, years later, it was found that although the risk of primary cardiac arrest was transiently increased during a *single* bout of vigorous exercise, *habitual* vigorous exercise was associated with an overall decrease in this risk (Siscovick *et al.*, 1984; Albert *et al.*, 2000). There are no absolute contraindications to very moderate exercise in chronic obstructive pulmonary disease patients (Pedersen and Saltin, 2006). However, in patients with asthma, a pause in training is recommended when an acute exacerbation occurs. In cases of infection, a pause in training is recommended until the patient has been asymptomatic for a day, where after training can be slowly resumed (Pedersen and Saltin, 2006).

Regarding muscle, bone and joint diseases, for instance, osteoarthritis and rheumatoid arthritis, exercise is contraindicated in cases of acute joint inflammation, if pain worsens after training and in cases of pericarditis and pleuritis (Pedersen and Saltin, 2006). The training of patients with osteoporosis should include activities with a low risk of falling (Pedersen and Saltin, 2006).

In cancer patients being treated with chemotherapy or radiotherapy, exercise is contraindicated when leukocyte concentrations fall below  $0.5 \times 10^9$  cells  $L^{-1}$ , haemoglobin below  $100 \text{ g} \cdot L^{-1}$ , thrombocyte concentration below  $20 \times 10^9$  cells  $L^{-1}$  and temperature above  $38^\circ\text{C}$ . Patients with bone metastases should not perform strength conditioning at high load. In cases of infection, a pause in training is recommended until the patient has been asymptomatic for a day, where after training can be slowly resumed (Pedersen and Saltin, 2006). A major concern is whether exercise training influences the anticancer effects of conventional cytotoxic therapy. The potential interaction between exercise and chemotherapy efficacy is biologically plausible. Indeed, earlier preclinical studies have reported both an inhibitory (Baracos, 1989) and augmentary (Thompson *et al.*, 1989) effect of endurance exercise training on mammary tumour growth and progression, although others have reported no association (Jones *et al.*, 2005).

In diabetic patients (both types I and II), exercise should be postponed if blood glucose is  $>2.5 \text{ g}\cdot\text{L}^{-1}$  together with ketonuria and  $>3.0 \text{ g}\cdot\text{L}^{-1}$  even without ketonuria, in both cases, before it is corrected. In patients with hypertension and active proliferative retinopathy, high-intensity training or training involving Valsalva-like manoeuvres should be avoided. Patients with neuropathy and incipient foot ulcers should refrain from activities entailing the bearing of the patient's own body weight.

In metabolic syndrome-related disorders, such as insulin resistance, dyslipaemia and obesity, there are no general contraindications; but training should take into account any comorbidities (Pedersen and Saltin, 2006). Finally, hypertensive patients with a blood pressure  $>180/105$  should begin pharmacotherapy before regular physical activity is initiated (relative contraindication) (Pescatello *et al.*, 2004). There is no evidence for an enhanced risk of sudden death or stroke in physically active persons with hypertension (Tipton, 1999). The American College of Sports Medicine (ACSM) recommends caution when performing very intensive dynamic exercise or strength conditioning with very heavy weights. Patients with left-sided cardiac hypertrophy should be particularly cautious about heavy strength conditioning. Patients with CHD should refrain from short intensive exercise situations.

It is well known that eccentric muscle contractions cause structural damage to muscle cells or inflammatory reactions within the muscles, as shown by an increase in the plasma activity of cytosolic enzymes and sarcolemma and Z-line disruption (Armstrong *et al.*, 1983). The severity of the damage and the extent of discomfort are exacerbated over time and can last for several days. The damaging effects of eccentric contractions can affect subsequent exercise sessions due to residual muscle pain, restriction of movement and reduced capacity to exercise at an intensity that may be beneficial for the exerciser (Howatson and van Someren, 2008). Thus, caution should be paid in exercise programs which include eccentric contractions especially in recreational or old practitioners.

## Exercise mimetics

Despite the clear evidence showing the powerful influence of exercise on health, physical inactivity remains a pressing public health issue. Technology and economic incentives tend to discourage activity: technology by reducing the energy needed for activities of daily living and economics by paying more for sedentary than for physically active work (Haskell *et al.*, 2007). Moreover, endurance exercise can also be unapproachable for most people in whom it might be impractical because of physical limitations or, as mentioned in the previous section, side effects. This fact stimulates the search for exercise mimetics (or 'exercise pills') that mimic exercise and, therefore, it has been the focus of important research over the past decades (Goodyear, 2008).

As mentioned in a previous section, exercise improves performance by activating several pathways that induce genetic changes, particularly in skeletal muscle, to increase aerobic metabolism and vascularisation, to ultimately enhance performance (Narkar *et al.*, 2011). AMPK is activated

by exercise and is essential for the exercise-mediated switch to aerobic myofibres in skeletal muscle (Wojtaszewski *et al.*, 2000). AMPK stimulates catabolic and suppresses anabolic pathways in an effort to restore cellular ATP levels and is activated robustly in skeletal muscle by acute as well as by chronic exercise (Winder *et al.*, 2006; Matsakas and Narkar, 2010). Recently, it was reported that 5-aminoimidazole-4-carboxamide-1- $\beta$ -D-ribofuranoside (AICAR) can mimic the effects of exercise by increasing GLUT-4 protein, hexokinase activity, resting glycogen content and muscle mitochondria (Narkar *et al.*, 2008). Mitochondrial biogenesis is activated by AMPK (Jorgensen *et al.*, 2007). This may be explained because AMPK is present in a transcriptional complex with PPAR- $\delta$ , where it can potentiate receptor activity via direct protein-protein interaction and/or by phosphorylating and activating coactivators such as PGC-1 $\alpha$  (Jager *et al.*, 2007). Puigserver *et al.* (1998) identified PGC-1 $\alpha$  as the master regulator of mitochondrial biogenesis and fuel homeostasis in mammalian tissues and Kovacs *et al.* (2005) observed that PGC-1 $\alpha$  mediates metabolic remodelling of skeletal myocytes, mimics exercise and reverses lipid-induced mitochondrial inefficiency. Thus, exercise training and oxidative fibre type are associated with increased mRNA expression of PGC-1 $\alpha$  (Lin *et al.*, 2002). Over-expression of PGC-1 $\alpha$  in mice induces dramatic changes in skeletal muscle such as increased mitochondrial biogenesis and fibre remodelling (Lin *et al.*, 2005). Moreover, PGC-1 $\alpha$  may be activated by the  $\text{Ca}^{2+}$ -signalling pathway involving both calcineurin and  $\text{Ca}^{2+}$ /calmodulin-dependent kinase and by p38 MAPK (Schiaffino *et al.*, 2007).

Recently, we have found an age-associated lack of reactivity of PGC-1 $\alpha$  in response to exercise (Derbre *et al.*, 2011), as aged rats had the same response as mice with genetic deletion (knockout) of PGC-1 $\alpha$ . Our results highlight the role of PGC-1 $\alpha$  in the loss of mitochondrial biogenesis associated with aging and point to this important transcriptional co-activator as a target for pharmacological interventions to prevent age-associated sarcopenia (Derbre *et al.*, 2011). Modulation of PGC-1 $\alpha$  levels in skeletal muscle is crucial for the prevention and treatment of age-related disorders (Sandri *et al.*, 2006). Accordingly, we recently proposed several pharmacological, non-hormonal, interventions to prevent the loss of muscle mass and sarcopenia such as PGC-1 $\alpha$  activators, angiotensin II receptor antagonists and allopurinol (Sanchis-Gomar *et al.*, 2011a).

On the other hand, activation or over-expression of the transcription factor PPAR- $\delta$  in muscle also results in an increase in mitochondrial biogenesis and in the proportion of oxidative muscle fibres (Narkar *et al.*, 2008). This results in increased running endurance and protection against diet-induced obesity and type 2 diabetes. The contrary is also true: muscle-specific knockout of PPAR- $\delta$  results in an age-dependent loss of oxidative muscle fibres, running endurance and insulin sensitivity (Schuler *et al.*, 2006). Thus, several potent and selective PPAR- $\delta$  agonists such as GW1516 have been also identified and proposed as mimetic drugs for endurance exercise (Narkar *et al.*, 2008).

Resveratrol has also been considered as an exercise mimetic. High doses of resveratrol improve endurance (Lagouge *et al.*, 2006). The deacetylase enzyme Sir2 and its mammalian homologue SIRT1 have been identified as its putative primary targets. Resveratrol also activates AMPK in



1 cells in culture, and it has been proposed for the prevention  
2 of mitochondrial dysfunction (Ungvari *et al.*, 2011) and of  
3 the wasting disorders associated with mechanical unloading  
4 (Momken *et al.*, 2011).

5 Remodelling of skeletal muscle by exercise is extremely  
6 complex. Many targets are available to mimic muscle adap-  
7 tations induced by exercise. However, exercise is linked to  
8 other multiple physiological adaptations that affect the vast  
9 majority of organs. It seems premature to conclude that all  
10 these molecules or substances are mimetics of exercise until  
11 the effects in other organs have been fully investigated. Appli-  
12 cability of these compounds at this point in time is limited.  
13 Studies aimed to find potential drugs that mimic exercise are  
14 now being performed.

## 15 Concluding remarks

16 Exercise is so beneficial for health that it should be considered  
17 as a drug. As for any other drug, dosing is very important.  
18 Otherwise, unfavourable side effects may occur. Some of the  
19 favourable effects of exercise apply to the general population.  
20 Prominent amongst these are its role in prevention of many  
21 diseases and in the promotion of healthy longevity (see  
22 Figure 2). But exercise can also be considered as treatment of  
23 established diseases. These include commonly occurring  
24 conditions such as depression, diabetes or cardiovascular  
25 diseases.  
26  
27

## 28 Acknowledgements

29 We thank Mrs Marilyn Noyes for her kind help in reviewing  
30 the manuscript. Our work is supported by grants SAF2008-  
31 00270, SAF2009-08334 from the Spanish Ministry of Educa-  
32 tion and Science; PROMETEO/2010/074 from the Conselleria  
33 de Educaci3n de la Generalitat Valenciana. ISCIII2006-  
34 RED13-027 from the 'Red Tem3tica de Investigaci3n Coop-  
35 erativa en Envejecimiento y Fragilidad (RETICEF)', EU  
36 Funded COSTB35 and DPS2008- 06968 from Spanish Minis-  
37 try of Innovation and Science. Our studies have also been co  
38 financed by FEDER funds from the European Union.  
39  
40

## 41 Conflict of interest

42 The authors declare that no conflict of interest exists.  
43  
44

## 45 References

46  
47  
48  
49 Albert CM, Mittleman MA, Chae CU, Lee IM, Hennekens CH,  
50 Manson JE (2000). Triggering of sudden death from cardiac causes  
51 by vigorous exertion. *N Engl J Med* 343: 1355–1361.  
52  
53 Armstrong RB, Ogilvie RW, Schwane JA (1983). Eccentric  
54 exercise-induced injury to rat skeletal muscle. *J Appl Physiol* 54:  
80–93.

Baracos VE (1989). Exercise inhibits progressive growth of the  
Morris hepatoma 7777 in male and female rats. *Can J Physiol  
Pharmacol* 67: 864–870. 55  
56  
57  
Benito B, Gay-Jordi G, Serrano-Mollar A, Guasch E, Shi Y, Tardif JC  
*et al.* (2011). Cardiac arrhythmogenic remodeling in a rat model of  
long-term intensive exercise training. *Circulation* 123: 13–22. 58  
59  
60  
Berchtold MW, Brinkmeier H, Muntener M (2000). Calcium ion in  
skeletal muscle: its crucial role for muscle function, plasticity, and  
disease. *Physiol Rev* 80: 1215–1265. 61  
62  
63  
Black A, Black MM, Gensini G (1975). Exertion and acute coronary  
artery injury. *Angiology* 26: 759–783. 64  
65  
Blair SN, Kohl HW 3rd, Paffenbarger RS Jr, Clark DG, Cooper KH,  
Gibbons LW (1989). Physical fitness and all-cause mortality. A  
prospective study of healthy men and women. *JAMA* 262:  
2395–2401. 66  
67  
68  
69  
Blair SN, LaMonte MJ, Nichaman MZ (2004). The evolution of  
physical activity recommendations: how much is enough? *Am J  
Clin Nutr* 79: 913S–920S. 70  
71  
72  
Boreham CA, Kennedy RA, Murphy MH, Tully M, Wallace WF,  
Young I (2005). Training effects of short bouts of stair climbing on  
cardiorespiratory fitness, blood lipids, and homocysteine in  
sedentary young women. *Br J Sports Med* 39: 590–593. 73  
74  
75  
76  
Borras C, Gomez-Cabrera MC, Vina J (2011). The dual role of p53:  
DNA protection and antioxidant. *Free Radic Res* 45: 643–652. 77  
78  
Bortz WM 2nd, Angwin P, Mefford IN, Boarder MR, Noyce N,  
Barchas JD (1981). Catecholamines, dopamine, and endorphin  
levels during extreme exercise. *N Engl J Med* 305: 466–467. 79  
80  
81  
Bouchard C, Shephard RJ, Stephens T (1994). Physical Activity,  
Fitness, and Health: International Proceedings and Consensus  
Statement: [Second International Consensus Symposium on  
Physical Activity, Fitness, and Health, Held May 5 to May 9, 1992,  
in Toronto, Canada]. Human Kinetics Publishers: Champaign, IL;  
Leeds. 82  
83  
84  
85  
86  
87  
Carr DB, Bullen BA, Skrinar GS, Arnold MA, Rosenblatt M,  
Beitins IZ *et al.* (1981). Physical conditioning facilitates the  
exercise-induced secretion of beta-endorphin and beta-lipotropin in  
women. *N Engl J Med* 305: 560–563. 88  
89  
90  
91  
Caspersen CJ, Powell KE, Christenson GM (1985). Physical activity,  
exercise, and physical fitness: definitions and distinctions for  
health-related research. *Public Health Rep* 100: 126–131. 92  
93  
94  
Church TS, Earnest CP, Skinner JS, Blair SN (2007). Effects of  
different doses of physical activity on cardiorespiratory fitness  
among sedentary, overweight or obese postmenopausal women  
with elevated blood pressure: a randomized controlled trial. *JAMA*  
297: 2081–2091. 95  
96  
97  
98  
99  
Czurko A, Hirase H, Csicsvari J, Buzsaki G (1999). Sustained  
activation of hippocampal pyramidal cells by 'space clamping' in a  
running wheel. *Eur J Neurosci* 11: 344–352. 100  
101  
102  
Derbre F, Gomez-Cabrera MC, Nascimento AL, Sanchis-Gomar F,  
Martinez-Bello VE, Tresguerres JA *et al.* (2011). Age associated low  
mitochondrial biogenesis may be explained by lack of response of  
PGC-1alpha to exercise training. *Age (Dordr)* ••: ••–••. 103  
104  
105  
106  
Dishman RK, Berthoud HR, Booth FW, Cotman CW, Edgerton VR,  
Fleshner MR *et al.* (2006). Neurobiology of exercise. *Obesity (Silver  
Spring)* 14: 345–356. 107  
108  
109  
Drezner JA (2008). Contemporary approaches to the identification  
of athletes at risk for sudden cardiac death. *Curr Opin Cardiol* 23:  
494–501. 110  
111  
112

1	Dunn AL, Trivedi MH, O'Neal HA (2001). Physical activity	Huang LE, Arany Z, Livingston DM, Bunn HF (1996). Activation of	59
2	dose-response effects on outcomes of depression and anxiety. <i>Med</i>	hypoxia-inducible transcription factor depends primarily upon	60
3	<i>Sci Sports Exerc</i> 6 (Suppl.): S587–S597. Discussion 609–610.	redox-sensitive stabilization of its alpha subunit. <i>J Biol Chem</i> 271:	61
4	Dunstan DW, Daly RM, Owen N, Jolley D, Vulikh E, Shaw J <i>et al.</i>	32253–32259.	62
5	(2005). Home-based resistance training is not sufficient to maintain	Jager S, Handschin CS, Pierre J, Spiegelman BM (2007).	63
6	improved glycemic control following supervised training in older	AMP-activated protein kinase (AMPK) action in skeletal muscle via	64
7	individuals with type 2 diabetes. <i>Diabetes Care</i> 28: 3–9.	direct phosphorylation of PGC-1alpha. <i>Proc Natl Acad Sci U S A</i>	65
8	Fox SM 3rd, Haskell WL (1968). Physical activity and the	104: 12017–12022.	66
9	prevention of coronary heart disease. <i>Bull N Y Acad Med</i> 44:	Ji LL, Gomez-Cabrera MC, Steinhafel N, Vina J (2004). Acute	67
10	950–967.	exercise activates nuclear factor (NF)-kappaB signaling pathway in	68
11	Freimuth M, Moniz S, Kim SR (2011). Clarifying exercise addiction:	rat skeletal muscle. <i>FASEB J</i> 18: 1499–1506.	69
12	differential diagnosis, co-occurring disorders, and phases of	Jones LW, Eves ND, Courneya KS, Chiu BK, Baracos VE, Hanson J	70
13	addiction. <i>Int J Environ Res Public Health</i> 8: 4069–4081.	<i>et al.</i> (2005). Effects of exercise training on antitumor efficacy of	71
14	Gomez-Cabrera MC, Borrás C, Pallardo FV, Sastre J, Ji LL, Vina J	doxorubicin in MDA-MB-231 breast cancer xenografts. <i>Clin Cancer</i>	72
15	(2005). Decreasing xanthine oxidase mediated oxidative stress	<i>Res</i> 11: 6695–6698.	73
16	prevents useful cellular adaptations to exercise in rats. <i>J Physiol</i> ••:	Jorgensen SB, Trebak JT, Viollet B, Schjerling P, Vaulont S,	74
17	•••••.	Wojtaszewski JF <i>et al.</i> (2007). Role of AMPKalpha2 in basal,	75
18	Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A,	training-, and AICAR-induced GLUT4, hexokinase II, and	76
19	Borrás C, Pallardo FV <i>et al.</i> (2008a). Oral administration of vitamin	mitochondrial protein expression in mouse muscle. <i>Am J Physiol</i>	77
20	C decreases muscle mitochondrial biogenesis and hampers	<i>Endocrinol Metab</i> 292: E331–E339.	78
21	training-induced adaptations in endurance performance. <i>Am J Clin</i>	Koves TR, Li P, An J, Akimoto T, Slentz D, Ilkayeva O <i>et al.</i> (2005).	79
22	<i>Nutr</i> 87: 142–149.	Peroxisome proliferator-activated receptor-gamma co-activator	80
23	Gomez-Cabrera MC, Domenech E, Vina J (2008b). Moderate	1alpha-mediated metabolic remodeling of skeletal myocytes mimics	81
24	exercise is an antioxidant: upregulation of antioxidant genes by	exercise training and reverses lipid-induced mitochondrial	82
25	training. <i>Free Radic Biol Med</i> 44: 126–131.	inefficiency. <i>J Biol Chem</i> 280: 33588–33598.	83
26	Gomez-Cabrera MC, Vina J, Ji LL (2009). Interplay of oxidants and	Kujala UM (2011). Born to be rich, physically active, fit and	84
27	antioxidants during exercise: implications for muscle health. <i>Phys</i>	healthy? <i>Scand J Med Sci Sports</i> 20: 367.	85
28	<i>Sportsmed</i> 37: 116–123.	Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C,	86
29	Goodyear LJ (2008). The exercise pill – too good to be true? <i>N Engl</i>	Daussin F <i>et al.</i> (2006). Resveratrol improves mitochondrial	87
30	<i>J Med</i> 359: 1842–1844.	function and protects against metabolic disease by activating SIRT1	88
31	Goodyear LJ, Chang PY, Sherwood DJ, Dufresne SD, Moller DE	and PGC-1alpha. <i>Cell</i> 127: 1109–1122.	89
32	(1996). Effects of exercise and insulin on mitogen-activated protein	Laurin D, Verreault R, Lindsay J, MacPherson K, Rockwood K	90
33	kinase signaling pathways in rat skeletal muscle. <i>Am J Physiol</i> 271	(2001). Physical activity and risk of cognitive impairment and	91
34	(2 Pt 1): E403–E408.	dementia in elderly persons. <i>Arch Neurol</i> 58: 498–504.	92
35	Grealy MA, Johnson DA, Rushton SK (1999). Improving cognitive	Lee DC, Artero EG, Sui X, Blair SN (2010). Mortality trends in the	93
36	function after brain injury: the use of exercise and virtual reality.	general population: the importance of cardiorespiratory fitness. <i>J</i>	94
37	<i>Arch Phys Med Rehabil</i> 80: 661–667.	<i>Psychopharmacol</i> 24 (4 Suppl.): 27–35.	95
38	Hakim AA, Petrovitch H, Burchfiel CM, Ross GW, Rodriguez BL,	Lee IM (2007). Dose-response relation between physical activity and	96
39	White LR <i>et al.</i> (1998). Effects of walking on mortality among	fitness: even a little is good; more is better. <i>JAMA</i> 297: 2137–2139.	97
40	nonsmoking retired men. <i>N Engl J Med</i> 338: 94–99.	Lee IM, Skerrett PJ (2001). Physical activity and all-cause mortality:	98
41	Harber VJ, Sutton JR (1984). Endorphins and exercise. <i>Sports Med</i>	what is the dose-response relation? <i>Med Sci Sports Exerc</i> 33 (6	99
42	1: 154–171.	Suppl.): S459–S471. Discussion S493–S454.	100
43	Hartley PH, Llewellyn GF (1939). Longevity of oarsmen. <i>Br Med J</i> 1:	Lejoyeux M, Avril M, Richoux C, Embouazza H, Nivoli F (2008).	101
44	657–662.	Prevalence of exercise dependence and other behavioral addictions	102
45	Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA <i>et al.</i>	among clients of a Parisian fitness room. <i>Compr Psychiatry</i> 49:	103
46	(2007). Physical activity and public health: updated	353–358.	104
47	recommendation for adults from the American College of Sports	Lin J, Wu H, Tarr PT, Zhang CY, Wu Z, Boss O <i>et al.</i> (2002).	105
48	Medicine and the American Heart Association. <i>Med Sci Sports Exerc</i>	Transcriptional co-activator PGC-1 alpha drives the formation of	106
49	39: 1423–1434.	slow-twitch muscle fibres. <i>Nature</i> 418: 797–801.	107
50	Holloszy JO, Coyle EF (1984). Adaptations of skeletal muscle to	Lin J, Handschin C, Spiegelman BM (2005). Metabolic control	108
51	endurance exercise and their metabolic consequences. <i>J Appl</i>	through the PGC-1 family of transcription coactivators. <i>Cell Metab</i>	109
52	<i>Physiol</i> 56: 831–838.	1: 361–370.	110
53	Howatson G, van Someren KA (2008). The prevention and	Lyons AS, Rjja P (1978). <i>Medicine: An Illustrated History</i> . H. N.	111
54	treatment of exercise-induced muscle damage. <i>Sports Med</i> 38:	Abrams: New York.	112
55	483–503.	Macera CA, Hootman JM, Sniezek JE (2003). Major public health	113
56	Hu FB, Willett WC, Li T, Stampfer MJ, Colditz GA, Manson JE	benefits of physical activity. <i>Arthritis Rheum</i> 49: 122–128.	114
57	(2004). Adiposity as compared with physical activity in predicting	Mandroukas K, Krotkiewski M, Hedberg M, Wroblewski Z,	115
58	mortality among women. <i>N Engl J Med</i> 351: 2694–2703.	Bjorntorp P, Grimby G (1984). Physical training in obese women.	116

1	Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO (2009).	Paffenbarger RS Jr, Brand RJ, Sholtz RI, Jung DL (1978). Energy	57
2	Sudden deaths in young competitive athletes: analysis of 1866	expenditure, cigarette smoking, and blood pressure level as related	58
3	deaths in the United States, 1980–2006. <i>Circulation</i> 119:	to death from specific diseases. <i>Am J Epidemiol</i> 108: 12–18.	59
4	1085–1092.		
5	Matsakas A, Narkar VA (2010). Endurance exercise mimetics in	Palomero J, Broome CS, Rasmussen P, Mohr M, Nielsen B, Nybo L	60
6	skeletal muscle. <i>Curr Sports Med Rep</i> 9: 227–232.	<i>et al.</i> (2008). Heat shock factor activation in human muscles	61
7		following a demanding intermittent exercise protocol is attenuated	62
8	McCloskey DP, Adamo DS, Anderson BJ (2001). Exercise increases	with hyperthermia. <i>Acta Physiol (Oxf)</i> 193: 79–88.	63
9	metabolic capacity in the motor cortex and striatum, but not in the		
10	hippocampus. <i>Brain Res</i> 891: 168–175.	Pedersen BK, Saltin B (2006). Evidence for prescribing exercise as	64
11	Mittleman MA, Maclure M, Tofler GH, Sherwood JB, Goldberg RJ,	therapy in chronic disease. <i>Scand J Med Sci Sports</i> 16 (Suppl. 1):	65
12	Muller JE (1993). Triggering of acute myocardial infarction by	3–63.	66
13	heavy physical exertion. Protection against triggering by regular		
14	exertion. Determinants of Myocardial Infarction Onset Study	Pescatello LS, Franklin BA, Fagard R, Farquhar WB, Kelley GA,	67
15	Investigators. <i>N Engl J Med</i> 329: 1677–1683.	Ray CA (2004). American College of Sports Medicine position	68
16		stand. Exercise and hypertension. <i>Med Sci Sports Exerc</i> 36:	69
17	Momken I, Stevens L, Bergouignan A, Desplanches D, Rudwill F,	533–553.	70
18	Chery I <i>et al.</i> (2011). Resveratrol prevents the wasting disorders of	Polednak AP (1972). Longevity and cause of death among Harvard	71
19	mechanical unloading by acting as a physical exercise mimetic in	College athletes and their classmates. <i>Geriatrics</i> 27: 53–64.	72
20	the rat. <i>FASEB J</i> ••: ••–••.		
21	Moorstein B (1968). Life expectancy of Ivy League rowing crews.	van Praag H, Christie BR, Sejnowski TJ, Gage FH (1999). Running	73
22	<i>JAMA</i> 205: 106.	enhances neurogenesis, learning, and long-term potentiation in	74
23		mice. <i>Proc Natl Acad Sci U S A</i> 96: 13427–13431.	75
24	Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S,	Prout C (1972). Life expectancy of college oarsmen. <i>JAMA</i> 220:	76
25	Lehar J <i>et al.</i> (2003). PGC-1 $\alpha$ -responsive genes involved in	1709–1711.	77
26	oxidative phosphorylation are coordinately downregulated in		
27	human diabetes. <i>Nat Genet</i> 34: 267–273.	Puigserver P, Wu Z, Park CW, Graves R, Wright M, Spiegelman BM	78
28		(1998). A cold-inducible coactivator of nuclear receptors linked to	79
29	Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW (1953a).	adaptive thermogenesis. <i>Cell</i> 92: 829–839.	80
30	Coronary heart-disease and physical activity of work. <i>Lancet</i> 265:		
31	1111–1120; concl.	Reid MB (2001). Invited Review: redox modulation of skeletal	81
32		muscle contraction: what we know and what we don't. <i>J Appl</i>	82
33	Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW (1953b).	<i>Physiol</i> 90: 724–731.	83
34	Coronary heart-disease and physical activity of work. <i>Lancet</i> 265:		
35	1053–1057; contd.	Ristow M, Zarse K, Oberbach A, Klötting N, Birringer M,	84
36		Kiehntopf M <i>et al.</i> (2009). Antioxidants prevent health-promoting	85
37	Murphy MH, Blair SN, Murtagh EM (2009). Accumulated versus	effects of physical exercise in humans. <i>Proc Natl Acad Sci U S A</i>	86
38	continuous exercise for health benefit: a review of empirical studies.	106: 8665–8670.	87
39	<i>Sports Med</i> 39: 29–43.		
40		Ruiz JR, Moran M, Arenas J, Lucia A (2010). Strenuous endurance	88
41	Myers J, Kaykha A, George S, Abella J, Zaheer N, Lear S <i>et al.</i> (2004).	exercise improves life expectancy: it's in our genes. <i>Br J Sports Med</i>	89
42	Fitness versus physical activity patterns in predicting mortality in	45: 159–161.	90
43	men. <i>Am J Med</i> 117: 912–918.		
44		Sakamoto K, Goodyear LJ (2002). Invited review: intracellular	91
45	Narkar VA, Downes M, Yu RT, Embler E, Wang YX, Banayo E <i>et al.</i>	signaling in contracting skeletal muscle. <i>J Appl Physiol</i> 93:	92
46	(2008). AMPK and PPAR $\delta$ agonists are exercise mimetics. <i>Cell</i>	369–383.	93
47	134: 405–415.		
48		Salminen A, Vihko V (1983). Endurance training reduces the	94
49	Narkar VA, Fan W, Downes M, Yu RT, Jonker JW, Alaynick WA <i>et al.</i>	susceptibility of mouse skeletal muscle to lipid peroxidation in	95
50	(2011). Exercise and PGC-1 $\alpha$ -independent synchronization of	vitro. <i>Acta Physiol Scand</i> 117: 109–113.	96
51	type I muscle metabolism and vasculature by ERR $\gamma$ . <i>Cell</i>		
52	<i>Metab</i> 13: 283–293.	Sanchis-Gomar F, Gomez-Cabrera MC, Vina J (2011a). The loss of	97
53		muscle mass and sarcopenia: non hormonal intervention. <i>Exp</i>	98
54	Neeper SA, Gomez-Pinilla F, Choi J, Cotman C (1995). Exercise and	<i>Gerontol</i> ••: ••–••.	99
55	brain neurotrophins. <i>Nature</i> 373: 109.		
56		Sanchis-Gomar F, Ollaso-Gonzalez G, Corella D,	100
		Gomez-Cabrera MC, Vina J (2011b). Increased average longevity	101
		among the 'Tour de France' cyclists. <i>Int J Sports Med</i> 32: 644–647.	102
		Sandri M, Lin J, Handschin C, Yang W, Arany ZP, Lecker SH <i>et al.</i>	103
		(2006). PGC-1 $\alpha$ protects skeletal muscle from atrophy by	104
		suppressing FoxO3 action and atrophy-specific gene transcription.	105
		<i>Proc Natl Acad Sci U S A</i> 103: 16260–16265.	106
		Schiaffino S, Sandri M, Murgia M (2007). Activity-dependent	107
		signaling pathways controlling muscle diversity and plasticity.	108
		<i>Physiology (Bethesda)</i> 22: 269–278.	109
		Schuler M, Ali F, Chambon C, Duteil D, Bornert JM, Tardivel A	110
		<i>et al.</i> (2006). PGC1 $\alpha$ expression is controlled in skeletal muscles	111
		by PPAR $\beta$ , whose ablation results in fiber-type switching, obesity,	112
		and type 2 diabetes. <i>Cell Metab</i> 4: 407–414.	113

1	Shephard RJ (2001). Absolute versus relative intensity of physical	Tipton CM (1999). Exercise training for the treatment of	35
2	activity in a dose-response context. <i>Med Sci Sports Exerc</i> 33 (6	hypertension: a review. <i>Clin J Sport Med</i> 9: 104.	36
3	Suppl.): S400–S418. Discussion S419–S420.		
4	Shephard RJ, Fitcher R (1997). Physical activity and cancer: how	Ungvari Z, Sonntag WE, de Cabo R, Baur JA, Csiszar A (2011).	37
5	may protection be maximized? <i>Crit Rev Oncog</i> 8: 219–272.	Mitochondrial protection by resveratrol. <i>Exerc Sport Sci Rev</i> 39:	38
6	Shirayama Y, Chen AC, Nakagawa S, Russell DS, Duman RS (2002).	128–132.	39
7	Brain-derived neurotrophic factor produces antidepressant effects in	Vaynman S, Gomez-Pinilla F (2006). Revenge of the ‘sit’: how	40
8	behavioral models of depression. <i>J Neurosci</i> 22: 3251–3261.	lifestyle impacts neuronal and cognitive health through molecular	41
9	Siscovick DS, Weiss NS, Fletcher RH, Lasky T (1984). The incidence	systems that interface energy metabolism with neuronal plasticity. <i>J</i>	42
10	of primary cardiac arrest during vigorous exercise. <i>N Engl J Med</i>	<i>Neurosci Res</i> 84: 699–715.	43
11	311: 874–877.		
12	Stuciak JA, Boylan C, Fritsche M, Altar CA, Lindsay RM (1996).	Vissing J, Andersen M, Diemer NH (1996). Exercise-induced	44
13	BDNF increases monoaminergic activity in rat brain following	changes in local cerebral glucose utilization in the rat. <i>J Cereb</i>	45
14	intracerebroventricular or intraparenchymal administration. <i>Brain</i>	<i>Blood Flow Metab</i> 16: 729–736.	46
15	<i>Res</i> 710: 11–20.		
16	Speed C, Jaques R (2010). High-performance sports medicine: an	Wannamethee SG, Shaper AG, Walker M (1998). Changes in	47
17	ancient but evolving field. <i>Br J Sports Med</i> 45: 81–83.	physical activity, mortality, and incidence of coronary heart disease	48
18	Sussman S, Sussman AN (2011). Considering the definition of	in older men. <i>Lancet</i> 351: 1603–1608.	49
19	addiction. <i>Int J Environ Res Public Health</i> 8: 4025–4038.	Warburton DE, Nicol CW, Bredin SS (2006a). Health benefits of	50
20	Teramoto M, Bungum TJ (2010). Mortality and longevity of elite	physical activity: the evidence. <i>CMAJ</i> 174: 801–809.	51
21	athletes. <i>J Sci Med Sport</i> 13: 410–416.	Warburton DE, Nicol CW, Bredin SS (2006b). Prescribing exercise as	52
22	Thompson HJ, Ronan AM, Ritacco KA, Tagliaferro AR (1989). Effect	preventive therapy. <i>CMAJ</i> 174: 961–974.	53
23	of type and amount of dietary fat on the enhancement of rat	Wen CP, Wai JP, Tsai MK, Yang YC, Cheng TY, Lee MC <i>et al.</i>	54
24	mammary tumorigenesis by exercise. <i>Cancer Res</i> 49: 1904–1908.	(2011). Minimum amount of physical activity for reduced mortality	55
25	Thompson PD, Crouse SF, Goodpaster B, Kelley D, Moyna N,	and extended life expectancy: a prospective cohort study. <i>Lancet</i>	56
26	Pescatello L (2001). The acute versus the chronic response to	•••••	57
27	exercise. <i>Med Sci Sports Exerc</i> 33 (6 Suppl.): S438–S445. Discussion	Westerlind KC (2003). Physical activity and cancer prevention –	58
28	S452–S433.	mechanisms. <i>Med Sci Sports Exerc</i> 35: 1834–1840.	59
29	Thompson PD, Franklin BA, Balady GJ, Blair SN, Corrado D,	Winder WW, Taylor EB, Thomson DM (2006). Role of	60
30	Estes NA 3rd <i>et al.</i> (2007). Exercise and acute cardiovascular events	AMP-activated protein kinase in the molecular adaptation to	61
31	placing the risks into perspective: a scientific statement from the	endurance exercise. <i>Med Sci Sports Exerc</i> 38: 1945–1949.	62
32	American Heart Association Council on Nutrition, Physical Activity,	Wojtaszewski JF, Nielsen P, Hansen BF, Richter EA, Kiens B (2000).	63
33	and Metabolism and the Council on Clinical Cardiology.	Isoform-specific and exercise intensity-dependent activation of	64
34	<i>Circulation</i> 115: 2358–2368.	5′-AMP-activated protein kinase in human skeletal muscle. <i>J Physiol</i>	65
		528 (Pt 1) 221–226.	66












<b>Toppan Best-set Premedia Limited</b>	
Journal Code: BPH	Proofreader: Elsie
Article No: 1970	Delivery date: 10 May 2012
Page Extent: 12	Copyeditor:

## AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.

Query References	Query	Remark
q1	AUTHOR: Please check if the affiliation address is correct.	
q2	AUTHOR: As per journal style, if there are only two authors for a Reference, both author names should be cited. So, the Reference citations have been amended throughout according to this style. Please check the whole text and confirm that they are correct.	
q3	AUTHOR Please note that BJP does not accept the dL (=100mL) as a unit of volume as it is not a SI unit. I have corrected this to give concentration per litre You may prefer to express as mg mL <sup>-1</sup> Please check.	
q4	AUTHOR: Please supply the volume number and page range for Reference Derbre <i>et al.</i> , 2011.	
q5	AUTHOR: Please supply the volume number and page range for Reference Gomez-Cabrera <i>et al.</i> , 2005.	
q6	AUTHOR: Please supply the volume number and page range for Reference Momken <i>et al.</i> , 2011.	
q7	AUTHOR: Please supply the volume number and page range for Reference Sanchis-Gomar <i>et al.</i> , 2011a.	
q8	AUTHOR: Please supply the volume number and page range for Reference Wen <i>et al.</i> 2011.	
q9	AUTHOR: '1978' . . . Please add the author names for this Reference.	
q10	AUTHOR: Pate <i>et al.</i> , 1995 has not been included in the Reference List, please supply full publication details.	
q11	<b>AUTHOR:</b> '1996' . . . Please add the author names for this Reference.	

# MARKED PROOF

## Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ <sup>Ⓢ</sup>
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ <sup>Ⓢ</sup>
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↙
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Υ or Υ under character e.g. Υ or Υ
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	ʹ or ʸ and/or ʹ or ʸ
Insert double quotation marks	(As above)	“ or ” and/or ” or ”
Insert hyphen	(As above)	⊥
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	┌┐	┌┐
Close up	linking ○ characters	○
Insert or substitute space between characters or words	/ through character or ∧ where required	Υ
Reduce space between characters or words		↑